

Deformed Bosons: Combinatorics of Normal Ordering

P. BLASIAK^{a,b}, A. HORZELA^b,

^a*Laboratoire de Physique Théorique des Liquides, Université Pierre et Marie Curie,
Tour 24 - 2e ét., 4 Pl.Jussieu, F 75252 Paris Cedex 05, France*

^b*H.Niewodniczański Institute of Nuclear Physics, Polish Academy of Sciences,
ul. Eliasza-Radzikowskiego 152, PL 31342 Krakow, Poland
e-mail: blasiak@lptl.jussieu.fr, andrzej.horzela@ifj.edu.pl*

K. A. PENSON^a AND A. I. SOLOMON^{a,c}

^a*Laboratoire de Physique Théorique des Liquides, Université Pierre et Marie Curie,
Tour 24 - 2e ét., 4 Pl.Jussieu, F 75252 Paris Cedex 05, France*

^c*The Open University, Physics and Astronomy Department,
Milton Keynes MK7 6AA, United Kingdom
email: penson@lptl.jussieu.fr, a.i.solomon@open.ac.uk*

We solve the normal ordering problem for $(A^\dagger A)^n$ where A and $A^\dagger$ are one mode deformed $([A, A^\dagger] = [N + 1] - [N])$ bosonic ladder operators. The solution generalizes results known for canonical bosons. It involves combinatorial polynomials in the number operator N for which the generating function and explicit expressions are found. Simple deformations provide examples of the method.

PACS: 02.10.Ox, 03.65.Fd, 05.30.Jp

Key words: boson normal ordering, combinatorics

1 Introduction

We consider the general deformation of the boson algebra [1] in the form

$$[A, N] = A, \quad [A^\dagger, N] = -A^\dagger, \quad [A, A^\dagger] = [N + 1] - [N]. \quad (1)$$

In the above A and $A^\dagger$ are annihilation and creation operators, respectively, while the number operator N counts particles. It is defined in the Fock basis as $N|n\rangle = n|n\rangle$ and commutes with $A^\dagger A$. Because of that in any representation of (1) N can be written in the form $A^\dagger A = [N]$, where $[N]$ denotes an arbitrary function of N , usually called the "box" function. For general considerations we do not assume any realization of the number operator N and we treat it as an independent element of the algebra. Moreover, we do not assume any particular form of the "box" function $[N]$. Special cases, like $so(3)$ or $so(2, 1)$ algebras, will provide examples that show how such a general approach simplifies if an algebra and its realization are chosen.

In this note we give the solution to the problem of the normal ordering of a monomial $(A^\dagger A)^n$ in deformed annihilation and creation operators. A classical result due to J. Katriel [2] is that for canonical bosons a and $a^\dagger$, *i.e.*, for the Heisenberg-Weyl algebra, we have

$$(a^\dagger a)^n = \sum_{k=1}^n \mathbb{S}_{n,k} (a^\dagger)^k a^k, \quad (\text{for canonical bosons}) \quad (2)$$

where $\mathbb{S}_{n,k}$ are the Stirling numbers of the second kind which count the number of partitions of an n element set into k subsets. Namely, (2) links the problem to combinatorics. In the case of deformed bosons we cannot express the monomial $(A^\dagger A)^n$ as the combination of normally ordered expressions in $A^\dagger$ and A only. This was found for q -deformed bosons some time ago, [3], and recently for the R-deformed Heisenberg algebra related to the Calogero model, [4]. The number operator N occurs in the final formulae because moving creation operators to the left cannot get rid of N if it is assumed to be an independent element of the algebra. It appears that in general we can look for a solution of the form

$$(A^\dagger A)^n = \sum_{k=1}^n \mathbb{P}_{n,k}(N) (A^\dagger)^k A^k, \quad (3)$$

where the coefficients $\mathbb{P}_{n,k}(N)$ are functions of the number operator N which depend on the box function $[N]$. Following [3] we will call them *operator-valued deformed Stirling numbers* or just *deformed Stirling numbers*. In the sequel we give a comprehensive analysis of this generalization. We give recurrences satisfied by $\mathbb{P}_{n,k}(N)$ of (3) and construct their generating functions. This will enable us to give $\mathbb{P}_{n,k}(N)$ explicitly and to demonstrate how the method works on examples of simple Lie-type deformations of the canonical case.

2 Recurrence relations

One may check by induction that for each $k \geq 1$ the following relation holds

$$[A^k, A^\dagger] = ([N + k] - [N])A^{k-1}. \quad (4)$$

Using this relation it is also easy to check by induction that deformed Stirling numbers satisfy the recurrences¹⁾

$$\mathbb{P}_{n+1,k}(N) = \mathbb{P}_{n,k-1}(N) + ([N] - [N - k])\mathbb{P}_{n,k}(N) \quad \text{for } 1 < k < n \quad (5)$$

with initial values

$$\mathbb{P}_{n,1}(N) = ([N] - [N - 1])^{n-1}, \quad \mathbb{P}_{n,n}(N) = 1. \quad (6)$$

The proof can be deduced from the equalities

$$\begin{aligned} \sum_{k=1}^{n+1} \mathbb{P}_{n+1,k}(N) (A^\dagger)^k A^k &= (A^\dagger A)^{n+1} = (A^\dagger A)^n A^\dagger A \\ &= \sum_{k=1}^n \mathbb{P}_{n,k}(N) (A^\dagger)^k A^k A^\dagger A \stackrel{(4)}{=} \sum_{k=1}^n \mathbb{P}_{n,k}(N) (A^\dagger)^k (A^\dagger A + [N + k] - [N]) A^k \\ &\stackrel{(1)}{=} \sum_{k=2}^{n+1} \mathbb{P}_{n,k-1}(N) (A^\dagger)^k A^k + \sum_{k=1}^n ([N] - [N - k])\mathbb{P}_{n,k}(N) (A^\dagger)^k A^k. \end{aligned}$$

¹⁾ The recurrence relation Eq.(5) holds for all n and k if one imposes the following "boundary conditions": $\mathbb{P}_{i,j}(N) = 0$ for $i = 0$ or $j = 0$ or $i < j$, except $\mathbb{P}_{0,0}(N) = 1$.

It can be shown that each $\mathbb{P}_{n,k}(N)$ is a homogenous polynomial of order $n - k$ in the variables $[N], \dots, [N - k]$. For a polynomial box function one gets $\mathbb{P}_{n,k}(N)$ as a polynomial in N . As we shall show, a simple example of the latter is the deformation of the $so(3)$ or $so(2, 1)$ Lie algebras.

Note that for the "box" function $[N] = N + \text{const}$ (*i.e.*, for the canonical algebra) we get the "ordinary" Stirling numbers [5] of the second kind which fulfill the recurrence relation $\mathbb{S}_{n+1,k} = \mathbb{S}_{n,k-1} + k \mathbb{S}_{n,k}$ with initial values $\mathbb{S}_{n,n} = \mathbb{S}_{n,1} = 1$.

3 Generating functions and general expressions

We define the set of ordinary generating functions (OGF) of polynomials $\mathbb{P}_{n,k}(N)$, for $k \geq 1$, in the form

$$P_k(x, N) := \sum_{n=k}^{\infty} \mathbb{P}_{n,k}(N) x^n. \quad (7)$$

The initial conditions (6) for $k = 1$ give

$$P_1(x, N) = \frac{x}{1 - ([N] - [N - 1]) x}. \quad (8)$$

Using the recurrences (5) supplemented by (6) one finds the relation

$$P_k(x, N) = \frac{x}{1 - ([N] - [N - k]) x} P_{k-1}(x, N), \quad \text{for } k > 1, \quad (9)$$

for which the proof is provided by the equalities

$$\begin{aligned} P_k(x, N) &\stackrel{(5)}{=} \sum_{n=k}^{\infty} (\mathbb{P}_{n-1,k-1}(N) + ([N] - [N - k]) \mathbb{P}_{n-1,k}(N)) x^n \\ &= x \sum_{n=k}^{\infty} \mathbb{P}_{n-1,k-1}(N) x^{n-1} + ([N] - [N - k]) x \sum_{n=k}^{\infty} \mathbb{P}_{n-1,k}(N) x^{n-1} \\ &= x P_{k-1}(x, N) + ([N] - [N - k]) x P_k(x, N). \end{aligned}$$

The expressions (8) and (9) give explicit formulae for the OGF

$$P_k(x, N) = \prod_{j=1}^k \frac{x}{1 - ([N] - [N - j]) x}. \quad (10)$$

For the canonical algebra this becomes in the OGF for Stirling numbers:

$$P_k(x) = \frac{x}{(1 - x)(1 - 2x) \cdots (1 - kx)}. \quad (11)$$

The explicit knowledge of the OGF (10) enables us to find the $\mathbb{P}_{n,k}(N)$ in a compact form. As a rational function of x it can be expressed as a sum of partial fractions

$$P_k(x, N) = x^k \sum_{r=1}^k \frac{\alpha_r}{1 - ([N] - [N - r]) x}, \quad (12)$$

where

$$\alpha_r = \frac{1}{\prod_{j=1, j \neq r}^k \left(1 - \frac{[N] - [N-j]}{[N] - [N-r]}\right)}. \quad (13)$$

From the definition (7) we have $\mathbb{P}_{n,k}(N) = [x^n] P_k(x, N)^1$. Expanding the fractions in the above equations and collecting the terms we get $\mathbb{P}_{n,k}(N) = \sum_{r=1}^k \alpha_r ([N] - [N-r])^{n-k}$. Finally, using (13), we arrive at

$$\mathbb{P}_{n,k}(N) = \sum_{r=1}^k \frac{([N] - [N-r])^{n-1}}{\prod_{j=1, j \neq r}^k ([N-j] - [N-r])} \quad (14)$$

where monotonicity of $[N]$ has been assumed. For the canonical case this yields the classical Stirling numbers $\mathbb{S}_{n,k} = \sum_{r=1}^k (-1)^{k-r} \frac{r^n}{r! (k-r)!}$, [5].

The first four families of general deformed Stirling polynomials may be obtained from (3) with a little help from standard computer algebra tools, and are

$$\mathbb{P}_{1,1}(N) = 1,$$

$$\begin{aligned} \mathbb{P}_{2,1}(N) &= [N] - [N-1], \\ \mathbb{P}_{2,2}(N) &= 1, \end{aligned}$$

$$\begin{aligned} \mathbb{P}_{3,1}(N) &= ([N] - [N-1])^2, \\ \mathbb{P}_{3,2}(N) &= 2[N] - [N-1] - [N-2], \\ \mathbb{P}_{3,3}(N) &= 1, \end{aligned}$$

$$\begin{aligned} \mathbb{P}_{4,1}(N) &= ([N] - [N-1])^3, \\ \mathbb{P}_{4,2}(N) &= 3[N]^2 - 3[N][N-1] - 3[N][N-2] + [N-1]^2 + [N-2][N-1] + [N-2]^2, \\ \mathbb{P}_{4,3}(N) &= 3[N] - [N-1] - [N-2] - [N-3], \\ \mathbb{P}_{4,4}(N) &= 1. \end{aligned}$$

4 Examples of simple deformations

If we fix the "box" function as $[N] = \mp \frac{N(N-1)}{2}$ then (1) gives the commutation relations of $so(3)$ and $so(2,1)$ algebras, respectively. The first four families of polynomials, satisfying $\mathbb{P}_{n,k}^{so(2,1)}(N) = (-1)^{n-k} \mathbb{P}_{n,k}^{so(3)}(N)$, because of (5) and (6),

¹⁾ This notation comes from combinatorics and denotes the coefficient multiplying $x^n/n!$ in the formal Taylor expansion of the generating function.

are

so(3)	so(2,1)
$\mathbb{P}_{1,1}(N) = 1$	$\mathbb{P}_{1,1}(N) = 1$
$\mathbb{P}_{2,1}(N) = -N + 1$	$\mathbb{P}_{2,1}(N) = N - 1$
$\mathbb{P}_{2,2}(N) = 1$	$\mathbb{P}_{2,2}(N) = 1$
$\mathbb{P}_{3,1}(N) = (N - 1)^2$	$\mathbb{P}_{3,1}(N) = (N - 1)^2$
$\mathbb{P}_{3,2}(N) = -3N + 4$	$\mathbb{P}_{3,2}(N) = 3N - 4$
$\mathbb{P}_{3,3}(N) = 1$	$\mathbb{P}_{3,3}(N) = 1$
$\mathbb{P}_{4,1}(N) = -(N - 1)^3$	$\mathbb{P}_{4,1}(N) = (N - 1)^3$
$\mathbb{P}_{4,2}(N) = 7N^2 - 19N + 13$	$\mathbb{P}_{4,2}(N) = 7N^2 - 19N + 13$
$\mathbb{P}_{4,3}(N) = -6N + 10$	$\mathbb{P}_{4,3}(N) = 6N - 10$
$\mathbb{P}_{4,4}(N) = 1$	$\mathbb{P}_{4,4}(N) = 1$

Up to now we have been considering A , $A^\dagger$, and N as independent elements of the algebra. Choosing a representation and expressing N in terms of A and $A^\dagger$ we arrive at expressions like (3) which can be normally ordered further. We may consider a realization of $so(2, 1)$ in terms of canonical operators a and $a^\dagger$

$$A = \frac{1}{2\sqrt{2}}aa \quad A^\dagger = \frac{1}{2\sqrt{2}}a^\dagger a^\dagger \quad N = \frac{1}{4} \{a^\dagger, a\}. \quad (15)$$

The solution of the normal ordering problem for $((a^\dagger)^2 a^2)^n$, [6], leads to

$$((a^\dagger)^2 a^2)^n = \sum_{k=2}^{2n} \mathbb{S}_{2,2}(n, k) a^{\dagger k} a^k, \quad (16)$$

where $\mathbb{S}_{2,2}(n, k)$ are generalized Stirling numbers of the second kind defined by

$$\begin{aligned} \mathbb{S}_{2,2}(n, k) &= \frac{(-1)^k}{k!} \sum_{p=2}^k (-1)^p \binom{k}{p} [p(p-1)]^n \\ &= \sum_{l=0}^n (-1)^l \binom{n}{l} \mathbb{S}_{1,1}(2n-l, k). \end{aligned} \quad (17)$$

The numbers $\mathbb{S}_{2,2}(n, k)$ also have a combinatorial interpretation which generalize partitions of a set (classical Stirling numbers). Consider $2n$ distinguishable objects anti-correlated in pairs. This can be realized by colouring them in such a way that there are exactly two objects of each colour, *i.e.* there are n anti-correlated (coloured) pairs. Now, numbers $\mathbb{S}_{2,2}(n, k)$ count the partitions of that set into k subsets with the restriction that anti-correlated pairs are divided among different subsets, *i.e.* all objects in a subset are of different colours. This means that we restrict partitions of a $2n$ set by requiring anti-correlation of pairs. This interpretation will be taken further elsewhere [7].

Terms in (16) with k even are directly expressible by A and $A^\dagger$. If k is odd then the factor $a^\dagger a$ may be moved to the left and with the convention $\mathbb{S}_{2,2}(n, k) = 0$ for $k > 2n$ we get

$$((a^\dagger)^2 a^2)^n = \sum_{k=1}^n (\mathbb{S}_{2,2}(n, 2k) + \mathbb{S}_{2,2}(n, 2k+1)(a^\dagger a - 2k)) a^{\dagger 2k} a^{2k}. \quad (18)$$

Finally, we get

$$(A^\dagger A)^n = \sum_{k=1}^n 8^{k-n} (\mathbb{S}_{2,2}(n, 2k) + \mathbb{S}_{2,2}(n, 2k+1)(2N - 2k - 1/2)) A^{\dagger k} A^k. \quad (19)$$

which shows that the general solution involving higher order polynomials in N can be simplified when a particular realization of the algebra is postulated.

5 Summary

We considered the problem of the normal ordering of monomials $(A^\dagger A)^n$ for generally deformed boson creation and annihilation operators. The general solution involves some combinatorial-like polynomials for which ordinary generating functions and explicit formulas can be given. It was shown that for particular realizations of the deformed algebra the solution may be further simplified.

P.B. wishes to thank the Polish Ministry of Scientific Research and Information Technology for support under Grant no: 1P03B 051 26.

References

- [1] A.I. Solomon: Phys. Lett. A **196** (1984) 29.
- [2] J. Katriel: Lett. Nuovo Cim. **10** (1974) 565.
- [3] J. Katriel and M. Kibler: J. Phys A **25** (1992) 2683.
J. Katriel: Phys. Lett. A **273** (2000) 159.
- [4] C. Burdick and O.Navratil: *Normal ordering for deformed Heisenberg algebra involving the reflection operator*, unpublished
- [5] L. Comtet: *Advanced Combinatorics*, Dordrecht: Reidel, 1974.
- [6] P. Blasiak, K. A. Penson, and A. I. Solomon: Phys. Lett. A **309** (2003) 198.
P. Blasiak, K. A. Penson, and A. I. Solomon: Ann. Combinatorics **7** (2003) 127.
- [7] P. Blasiak, A. Horzela, K. A. Penson, and A. I. Solomon: to be published.